

Scotland's Rural College

## Causes of keel bone damage and their solutions in laying hens

Harlander-Matauschek, A; Rodenburg, TB; Sandilands, V; Tobalske, BW; Toscano, MJ

*Published in:*  
World's Poultry Science Journal

*DOI:*  
[10.1017/S0043933915002135](https://doi.org/10.1017/S0043933915002135)

Print publication: 01/09/2015

*Document Version*  
Peer reviewed version

[Link to publication](#)

*Citation for pulished version (APA):*  
Harlander-Matauschek, A., Rodenburg, TB., Sandilands, V., Tobalske, BW., & Toscano, MJ. (2015). Causes of keel bone damage and their solutions in laying hens. *World's Poultry Science Journal*, 71(3), 461 - 472.  
<https://doi.org/10.1017/S0043933915002135>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Abbreviated title**

Keel Bone Damage: Causes and solutions

**Summary (100-300 words)**

Keel bone damage (KBD) is a critical issue facing the contemporary laying hen industry due to the likely pain leading to compromised welfare and reduced productivity. Recent reports suggest that KBD, while highly variable and likely dependent on a host of factors, extends to all housing systems (including traditional battery cages, furnished cages and non-cage systems), genetic lines, and management styles. Despite the extent of the problem, the research community remains uncertain as to the causes and influencing factors of KBD. To combat these issues, the current review was produced following discussions from the 1<sup>st</sup> International Keel Bone Damage Workshop held in Switzerland in April 2014. This endeavor sought to assess current knowledge, foster novel collaborations, propose unique methodologies and highlight the key areas where innovative research is needed. The current paper is based on the content of those discussions and presents nine recommendations for future research efforts.

**Keywords: Keel, fracture, damage, welfare, bone, laying hen**

**Introduction**

The high frequency of damage seen in the keel bone (a bone extending from the sternum) of laying hens within commercial systems represents one of the greatest welfare problems facing the industry as suggested by the UK's Farm Animal Welfare Committee (FAWC, 2010, 2013). Beyond the obvious welfare issue of gross skeletal deformity, concern stems from the likely associated pain indicated by the decreased latency to descend from a perch (Nasr *et al.*, 2012a; c, 2014). One type of KBD, keel fractures (KF), also referred to as old breaks, can be defined as breaks in the bone that will typically manifest as a callus around the fracture site after

a few days; KF may also involve sharp, unnatural deviations, or bending, of the bone (Wilkins *et al.*, 2004). The problem of keel damage is clearly widespread with similar yet highly variable levels reported in various countries (such as: Switzerland (Kappeli *et al.*, 2011); UK (Wilkins *et al.*, 2011); The Netherlands, Belgium, Germany (Rodenburg *et al.*, 2008; Heerkens *et al.*, 2013); and Canada (Petrik *et al.*, 2014)). Keel bone damage (KBD) extends across genetic lines (Kappeli *et al.*, 2011) and all types of housing systems (Wilkins *et al.*, 2011; Petrik *et al.*, 2015) including organic production systems (Bestman and Wagenaar, 2014). Despite their ubiquity, the causes and influencing factors of KBD remain largely unknown to the research community, a circumstance that severely handicaps the development of effective strategies to reduce their occurrence and severity.

To identify critical areas where research is needed and coordinate the multiple ongoing and future research efforts of laboratories, the International Keel Bone Damage Workshop was organized by the University of Bern in April 2014. This manuscript serves to highlight the workshop dialogue and harness the collective energies of the research community towards reducing KBD in laying hens by formulating key recommendations for forthcoming research.

**Recommendation 1 – A uniform methods of evaluating KBD should be employed with relevant criteria to ensure reliability of results.**

Currently, evaluating KBD in live birds is principally done by palpation; a method that originated primarily from two papers (Wilkins *et al.*, 2004; Scholz *et al.*, 2008) which have since been adopted by a variety of labs. While broadly similar in their classification of damage, key differences exist and require resolution. Most critically, Wilkins *et al.* (2004) only discussed KF while Scholz *et al.* (2008) included deviations of the keel (deformations from a theoretically flat, two dimensional plane; also referred to as curving, bending, etc.) as well. Given that KF and deviations likely result from different causal factors (i.e., sudden impacts

causing large forces vs. sustained and small forces, respectively), this lack of clarity represents a major obstacle in combining results to allow for comprehensive interpretation of the problem. The multiple methods in use are a challenge in resolving KBD, hindering our abilities to compare findings and resulting in needless and inefficient replication. We recommend development of a uniform assessment of KBD that assesses KF and deviations where each uses a binary scale of whether damage is present. The system should be developed in concert with an associated scoring sheet which should be made publicly available to facilitate the amalgamation of data produced by the labs conducting related research.

Beyond defining the presence or absence of damage, identifying grades of severity is also needed to determine the types of KBD that are a concern. Research that can provide reliable and consistent markers to classify KBD in grades of severity that are grounded in meaningful criteria relevant to animal welfare (e.g., pain caused by fracture) and/or other spheres of concern (e.g. productivity) is essential.

We also recommend adoption of training criteria for persons assessing KBD (particularly by palpation which is more subjective than dissection) to ensure greater reliability of results. A system employed by assessors with appropriate training can produce results that are more useful in determining true occurrence of damage and evaluating interventions. The method of assessment, including free access to a developed scoring form, the described definitions for the various forms of damage, as well as criteria and training for palpation, is described elsewhere (Casey-Trott *et al.*, *submitted to Poultry Science*).

## **Recommendation 2 – Investigate low energy, non-collision events as a source of KF.**

High energy events within non-cage systems provide a likely mechanism for KF that have been suggested to result from collisions with elements of animal housing such as perches (Scott *et al.*, 1997; Moinard *et al.*, 2004a; Sandilands *et al.*, 2009; Wilkins *et al.*, 2011). Counter-

intuitively, KF also frequently occur in cage systems where there exists seemingly limited opportunities for dynamic loading as might occur during collision. Thus, improving our understanding of the sources of KF will require insight of dynamic as well as static events. The mechanisms of KF in cage systems are likely not exclusive to this environment, i.e., static mechanisms are likely to cause damage in non-cage systems as well, and thus deserve investigation to provide a more comprehensive understanding of the sources of KBD.

Although commercial laying hens kept in enriched cages and non-cage systems have opportunities for weight-bearing activities, birds' skeletons remain fragile because of a combination of nutritional, environmental and genetic factors (Whitehead, 2004a). It would be worthwhile to test whether seemingly innocuous, quasi-static activities can induce KF. While perching, hens place considerable forces on their keels which is in contact with the perch (Pickel *et al.*, 2011) and certain perch designs or material may result in severe keel bone deviations and perhaps also KF (Pickel *et al.*, 2010). This mechanism would be similar to compression fractures in osteoporotic bones of humans, which have been shown to occur spontaneously or with normally innocuous activities, such as sneezing or twisting (Kondo, 2008). Furthermore, application of modeling techniques, including finite-element models (Tomaszewski *et al.*, 2010) that represent the keel bone, would be useful to provide an understanding of the static and dynamic loading patterns in the bone.

More generally, a broader understanding of the keel, using a comparison of relevant bird phylogenies, could be helpful to establish the morphological capacity of the keel and how modern housing conditions exceed that capacity. It would be useful to assess keel integrity in relation to the functional morphology of diverse avian clades that exhibit significant variation in hindlimb/forelimb modules, keel morphology, flight style, and use of natural perches (Heers and Dial, 2012).

**Recommendation 3 – Investigate the relationship between deviations and KF.**

A keel with deviations may lead to unequal bone loading during wing-flapping and concentration of strain energy in ways that increase the risk of fracture. The paired *pectoralis* muscles are capable of generating enormous force, work and power output during normal flight (Tobalske *et al.*, 2003) and these outputs increase dramatically during vigorous wing-flapping (Tobalske and Dial, 2000; Jackson and Dial, 2011). Comparable wing-flapping is often observed in commercial laying hens that slip from a perch or aviary tier and try to regain their footing (personal observation, M Toscano). Three-dimensional force-balance calculations (Hutchinson *et al.*, 2005; Baier *et al.*, 2006) should be used to test for effects of keel deformity upon bone loading during *pectoralis* muscle contraction; particularly those observed during episodes where balance is lost or panics (see Recommendation #5). Also, deviated keels may lead to KF indirectly by complicating balance maneuvers, an additional topic which deserves investigation.

**Recommendation 4 – Investigate the role of bird development in KBD susceptibility.**

It is of prime importance to learn more about development of locomotor and cognitive skills as these contribute towards the ability of hens to navigate within the home system. It is doubtful that the traditional aviary rearing system, where chicks are kept confined to platforms for the first four weeks after which the sides are opened, is the best system to prepare birds for aviary housing (Kozak *et al.* 2015). Under natural conditions, locomotor capacity is critical in Galliformes as they are confronted with immediate challenges to escape predators, search for food and seek shelter (Dial and Jackson, 2011). Beginning at six days post-hatching, wild Galliformes will readily flap their wings to produce aerodynamic forces that enhance hindlimb function while moving up inclines, a behaviour called wing-assisted incline running (WAIR) (Dial, 2003; Tobalske and Dial, 2007; Dial *et al.*, 2008). Adult Galliformes may also prefer WAIR rather than flight to reach an elevated area in a complex, natural habitat as well (Dial and Jackson, 2011). A

better understanding of this process in commercial strains could be helpful to design juvenile and/or adult hen housing systems that improve locomotor abilities (Le Blanc et al. 2015). For instance, variable-engineered systems would be suitable for accommodating the birds as they develop, using more ramps and adjustable angles with increasing age. A potential advantage of WAIR compared with flight may be that whole-body kinetic energy is less during WAIR (Tobalske and Dial, 2000, 2007), a benefit that could reduce the risk of KBD due to accidental impact with the housing environment. Additionally, increased wing-flapping during development may assist in improving balancing abilities (Filipa *et al.* 2010) while greater activity is known to correlate with bone strength (Rath *et al.*, 2000). Beyond musculo-skeletal development, juvenile birds using WAIR will likely learn neuromuscular coordination that will be useful for negotiating three-dimensional structures in adulthood, an impairment suggested by Gunnarsson *et al.* (2000). Research in this area will help to identify optimum rearing conditions likely to protect birds from cognitive impairment in adulthood.

**Recommendation 5 - Investigate the role of escape reactions as a source of KF.**

Efforts should also be made to understand the damage resulting from sudden escape reactions, or panics. Escape is normally triggered by a situation that is, or is perceived to be, life-threatening and thus is not “normal” in the context of commercial poultry husbandry, though has been documented (Richards *et al.*, 2012). Escapes resemble the pattern seen in Galliformes involving take-off using high-frequency, high-amplitude wing beats that feature enormous power output (Tobalske and Dial, 2000; Tobalske *et al.*, 2003), a quick return to the ground using a glide, and then resumption of walking or running. Given that the response is one of last resort, its manifestation is likely to exceed the keel’s morphological capacity. Additionally, escape flights will not allow for the precise navigation required in housing systems. A more thorough

understanding of the causes of escapes is necessary, including the roles of genetic selection and stockmanship, and the effects of dim lighting conditions and intra-bird spacing (Tillmann, 2009).

A comparative evolutionary approach could further aid investigation regarding the role of escape reactions as a source of damage. A museum survey of wild birds of different species showed that 4.5% out of a sample size of 6,212 specimens had sustained and survived bone injuries of which clavicle injuries were the greatest in number, especially in smaller birds, and were attributed to collisions with solid objects (Tiemeier, 1941). The sample included 45 birds in the Phasianidae, the family that includes the chicken, with an incidence rate of 10% in this family. These percentages are far lower than incidence rates of KBD in layer hens, but they do suggest that the escape flight of birds in the Phasianidae (Tobalske and Dial, 2000) may be correlated with bone damage. An alternative explanation, however, is that some damage reported by Tiemeier (1941) was due to gunshot, as many species in the Phasianidae are gamebirds.

#### **Recommendation 6 - Investigate genetic capacity to reduce KBD.**

The role of genetics as a contributing factor in KBD dates back to work by Hyre (1955) who showed that the tendency to develop keel deformities was heritable by successfully selecting for and against KBD over six generations. Even earlier, Warren (1937) showed that crooked keel bones (in comparison to straight keels) had a reduced ash content which he suggested was a causal factor. In considering the scope for genetic selection against KBD, we must first determine what traits should be selected where possibilities include: stronger bones, improved physical ability, and increased docility. Alternatively, as KBD is a relatively complex trait with a number of genetic and environmental factors playing a role, genomic selection should be considered (Fulton, 2012) by carefully monitoring the incidence of KBD in a large population of laying hens and then comparing genomic information of hens with and hens without KBD. One of the strengths of this approach is that it does not target a single factor, e.g.



bone strength, but focuses on the actual presence or absence of damage. In theory, the associated mechanism(s) is (are) selected in the process of identifying birds with the desired trait. A challenge with the genomic approach is that it requires a large sampling population (e.g., >5,000) of which a clear KBD phenotype is needed. However, once the genomic fingerprint of a hen with no KBD is acquired, no further phenotypic measurements are required for the selection program (Eggen, 2012). Although breeding for reduced KBD is attractive, one should be aware of possible linkages and trade-offs with other traits, e.g., reduced egg shell thickness and egg breaking strength (Stratmann *et al.*, *in prep*). Whitehead (2004b) provides an excellent review of the relationship between skeletal integrity and egg quality.

**Recommendation 7 – Investigate housing adaptations that affect frequency of KBD.**

Large differences between housing systems in the incidence KF indicate that housing design and/or management plays a key-role (Rodenburg *et al.*, 2008; Wilkins *et al.*, 2011). Perches have received particular attention. As an indication that perches have a causal role in the occurrence of KBD, higher rates were reported at end of lay for hens in conventional cages with (92%) compared to those without (83%) metal perches (Hester *et al.*, 2013). Similarly, Wilkins *et al.*, (2011) reported a 10-34% increase in KF when perches were added in an organic mobile system. Often, round metal perches are used, which may not offer hens adequate support for their grip. Perches with slightly larger diameters and those made from more flexible materials (wood, rubber) have been suggested to be more capable of absorbing forces during impact and preventing KBD (Pickel *et al.*, 2010, 2011). Perches covered with a soft rubber layer were successful in reducing the number of keel fractures within a commercial aviary suggesting the benefit of this option (Stratmann *et al.*, Accepted to PlosONE), possibly not only by reducing the pressure on the keel, but also by providing a cushion and improved grip when landing (Scholz *et al.*, 2014). Perches as a source of fracture is discussed in more detail by Sandilands *et al.* (2009).

Apart from perches, the three-dimensional environment of the hens has to be designed so that it allows the hens to navigate between the different parts of the system. In most commercial aviary systems, improvements are possible that would improve hens' possibilities for safely navigating through the system. One option is to add ramps to aid transition between tiers. Stratmann *et al.* (2015) showed that adding ramps to a commercial aviary system reduced falls by 55%, collisions by 41% and keel fractures by 24%, while movements between tiers increased by 44%. The width of the corridors between the different rows within systems also needs further attention. If the corridor width is at the limit of the birds' navigational ability, it may cause increased collisions due to misjudged jumps (Heerkens *et al.*, 2014).

Another factor to be considered when determining optimum housing is lighting. In many commercial laying hen operations, light intensity is kept at a relatively low level, especially in flocks that are prone to develop feather pecking. Birds need sufficient light and contrast to make an appropriate jump and safe landing (Moinard *et al.*, 2004b). The timing of the light transition also seems important: Stratmann *et al.* (2013) reported that vertical movements occurred mainly during a dusk phase when hens usually move to a perching spot for the night. A sudden switch from light to dark periods may increase the risk of KF during this time. Hence, a gradual dawn- and dusk should be investigated as a possible means of reducing KF. Lastly, Heerkens *et al.* (2014) found flooring type to also be a factor where wire flooring had greater frequency of KF compared with plastic flooring, though the underlying causes need to be explored.

#### **Recommendation 8 – Investigate nutritional solutions to reduce KBD.**

Solutions should also include changes in bird management and nutrition. The high incidence of KF in cages (Hester *et al.*, 2013) could point to a calcium shortage. Egg shell formation takes place during the night and hens need a large amount of calcium at this time to produce an eggshell each day. Hens can mobilize this calcium partly from their bones, but they

need to restore their supply, otherwise the risk osteoporosis (Whitehead and Fleming, 2000). One way to supply hens with calcium during the night is to give them daytime access to calcium sources with larger particle sizes, such as grit or shells, which will then be digested during the night. Larger particle sizes of calcium has been shown to benefit skeletal health (Cheng and Coon, 1990; Guinotte *et al.*, 1995) including that of the keel (Fleming *et al.*, 1998), though usage of this technique varies due to multiple factors including damage to feeding equipment and birds selectively eating the larger particles. Thus, there may be a benefit in supplying grit separately, or developing other sources of calcium that help the hens to restore their supplies and prevent bone weakness. Other nutritional changes could involve incorporation of omega-3 content into the diet which has been shown to result in reduced fracture incidence (Toscano *et al.*, Accepted to Poultry Science; Tarlton *et al.*, 2013) possibly by modulating bone metabolism and modeling (Liu *et al.*, 2003; Watkins *et al.*, 2003; Baird *et al.*, 2008).

#### **Recommendation 9 – Investigate and quantify KBD and production losses**

Physical conditions that are associated with pain can, if severe enough, induce redistribution of endogenous resources and derail physiological processes that ensure long-term survival, a classical criterion for compromised welfare (Moberg, 1985; Broom, 1991). More specifically, Prunier *et al.*, (2013) advocated changes in productivity as an indication of pain and potentially for compromised welfare. As mentioned before, recent work has shown that individual birds with fractures housed in large groups (~350 birds/group) produced eggs that were characterized with reduced breaking strength and thinner shells (Toscano, *in prep*). This response may represent a diversion of resources where minerals (e.g. calcium) and energy, normally directed towards egg production, must consequently be reallocated to the process of healing bone (Thiruvankadan *et al.*, 2010). Similar results for altered egg production in individual birds were found by Nasr *et al.* (2012b, 2013), although this was assessed in non-commercial conditions

(i.e., individual hens isolated in separate cages) in order to link the egg and hen. More critically, work by Nasr *et al.* (2012b, 2013), as well as that by Toscano (*in prep*), did not control for natural variation in bird laying capacity. The lack of pre-KF data leaves open the possibility that birds prone to KF may produce less and weaker eggs independent of whether KFs occurred. Differences in egg character after fracture must be shown to be absent beforehand if the measure is to be a valid indicator of welfare. Interestingly, others have been unable to demonstrate a link between egg production and keel fracture at the flock level (Heerkens *et al.*, 2013). The lack of a relationship could be due to high flock-level variance rather than the absence of an effect, indicating the need for research at bird level responses. Alternatively, Whitehead (2004b), reviewing several studies that examined individually housed birds bred for different bone qualities, suggested that little correlation existed between egg production and bone quality. The finding was supported by Gebhardt-Henrich and Fröhlich (2012) who reported more fractures in hens which laid their first egg earlier. As an additional complication in linking the occurrence of fractures with production data, the period in which fractures are seen to most dramatically increase (25 -35 weeks of age) is also the one in which birds are coming off peak of lay, thus a drop in production is expected independent of fractures. Therefore, the predicted falloff in egg production resulting from KBD may be subsumed by the drop in egg production as the hen exits the peak of lay period.

As discussed above, quantification of production endpoints and the loss of productivity associated with KBD can be used as a powerful means to assess changes in animal welfare. More so, because concern for animal welfare is not globally consistent (Lopez, 2007) with the strongest interest in Europe and North America, framing the problems of KBD in terms of productivity losses and compromised profit could provide alternative motivations that move towards an ultimate goal of reducing KBD. This particular argument is powerful as it does not

diminish the reality that action on the grounds of compromised welfare is necessary, but rather adds a supplementary dimension that will drive stakeholders to effect change.

Interestingly, it is often suggested that KF result from bone that is weakened by the process of demineralization to provide adequate amounts of calcium for egg shell formation. If correct, continued egg production should associate with a decrease in bone strength and an increase in the occurrence of KF. While this appears to be the case for the first 20 weeks of egg production, recent comparisons of several studies suggest that rates of fractures actually appear to flatten and possibly fall after 45 wks of age (Stratmann et al., accepted to Applied Animal Behaviour Science; Toscano et al., accepted to Poultry Science; Tarlton et al., 2013; Petrik et al., 2014). It is possible that this decrease could be attributed to altered behaviour, though use of an *ex vivo* impact testing protocol with dead hens (Toscano *et al.*, 2013) identified a pattern of decreased susceptibility to fracture (Toscano *et al.*, 2014) that mirrored the on-farm observations of live hens. Further research is needed to determine how this change in fracture occurrence relates to altered bone physiology and egg production during this period.

## **Overall Conclusions**

Keel bone damage represents a welfare and productivity problem for the laying hen industry and, while achievements have been made in understanding the nature and cause of occurrence, we remain far from resolving the issue. The current paper highlights areas of research that would achieve the goal of reducing KBD, encourage adoption of methods to improve the accuracy and reliability of reporting, and provide technical changes that could be adopted.

## **References**

**BAIER, D. B., GATESY, S. M. and JENKINS, F. A.** (2006) A critical ligamentous mechanism in the evolution of avian flight. *Nature* Vol. **445**: 307–310.

288    **BAIRD, H. T., EGGETT, D. L. and FULLMER, S.** (2008) Varying ratios of omega-6: omega-  
289    3 fatty acids on the pre-and postmortem bone mineral density, bone ash, and bone breaking  
290    strength of laying chickens. *Poultry Science* **87**: 323–328

291    **BESTMAN, M. and WAGENAAR, J. P.** (2014) Health and Welfare in Dutch Organic Laying  
292    Hens. *Animals* **4**: 374–390.

293    **BROOM, D. M.** (1991) Animal Welfare: Concepts and measurements. *Journal of Animal*  
294    *Science* **69**: 4167–4175.

295    **CHENG, T. K. and COON, C. N.** (1990) Effect of calcium source, particle size, limestone  
296    solubility in vitro, and calcium intake level on layer bone status and performance. *Poultry*  
297    *Science* **69**: 2214–2219.

298    **DIAL, K. P.** (2003) Wing-assisted incline running and the evolution of flight. *Science* **299**: 402–  
299    404.

300    **DIAL, K. P. and JACKSON, B. E.** (2011) When hatchlings outperform adults: locomotor  
301    development in Australian brush turkeys (*Alectura lathami*, Galliformes). *Proceedings Royal*  
302    *Society Biological Sciences* **278**: 1610–1616.

303    **DIAL, K. P., JACKSON, B. E. and SEGRE, P.** (2008) A fundamental avian wing-stroke  
304    provides a new perspective on the evolution of flight. *Nature* **451**: 985–989.

305    **EGGEN, A.** (2012) The development and application of genomic selection as a new breeding  
306    paradigm. *Animal Frontiers* **2**: 10–15.

307    **FAWC.** (2010) Opinion on Osteoporosis and Bone Fractures in Laying Hens . *Farm Animal*  
308    *Welfare Council*, London.

309    **FAWC.** (2013) An open letter to Great Britain Governments: Keel bone fracture in laying hens.

310 **FILIPA, A., BYRNES, R., PATERNO, M.V., MYER, G.D., AND HEWETT., T.E.** (2010)  
 311 Neuromuscular training improves performance on the star excursion balance test in young female  
 312 athletes. *Journal of Orthopaedic and Sports Physical Therapy* **40**: 551-558.  
 313 **FLEMING, R. H., MCCORMACK, H. A. and WHITEHEAD, C. C.** (1998) Bone structure  
 314 and strength at different ages in laying hens and effects of dietary particulate limestone, vitamin  
 315 K and ascorbic acid. *British Poultry Science* **39**: 434–440.  
 316 **FULTON, J. E.** (2012) Genomic selection for poultry breeding. *Animal Frontiers* **2**: 30–36.  
 317 **GEBHARDT-HENRICH, S. and FRÖLICH, E. K. F.** (2012) Auftreten von  
 318 Brustbeinfrakturen und individuelles Verhalten bei Legehennen. ERHARD, M., POLLMAN, U.,  
 319 PUPPE, B., REITER, K. and WAIBLINGER, S. (Eds) *KTBL*, pp. 52-60 (Freiburg, Germany).  
 320 **GUINOTTE, F., GAUTRON, J., NYS, Y. and SOURMARMON, A.** (1995) Calcium  
 321 solubilization and retention in the gastrointestinal tract in chicks (*Gallus domesticus*) as a  
 322 function of gastric acid secretion inhibition and of calcium carbonate particle size. *British*  
 323 *Journal of Nutrition* **73**: 125–139.  
 324 **GUNNARSSON, S., YNGVESSON, J., KEELING, L. J. and FORKMAN, B.** (2000) Rearing  
 325 without early access to perches impairs the spatial skills of laying hens. *Applied Animal*  
 326 *Behaviour Science* **67**: 217–228.  
 327 **HEERKENS, J., DELEZIE, E., KEMPEN, I., ZOONS, J., RODENBURG, T. B. and**  
 328 **TUYTENS, F.** (2013) Do keel bone deformations affect egg-production in end-of-lay housing  
 329 hens housed in aviaries? TAUSON, R., BLOKHUIS, H. J., BERG, L., ELSON, A. (Eds) *9th*  
 330 *European Poultry Conference*, pp.127 (Uppsala, Sweden).  
 331 **HEERKENS, J. L. T., KEMPEN, I., ZOONS, J., DELEZIE, E., RODENBURG, T. B.,**  
 332 **AMPE, B. and TUYTENS, F. A. M.** (2014) Effect of aviary housing characteristics on laying

333 hen welfare and performance. *Proceedings of the 48th Congress of the International Society for*  
334 *Applied Ethology*, Vitoria-Gasteiz, Spain pp. 158.

335 **HEERS, A. M. and DIAL, K. P.** (2012) From extant to extinct: locomotor ontogeny and the  
336 evolution of avian flight. *Trends in Ecology. and Evolution.* **27**: 296–305.

337 **HESTER, P. Y., ENNEKING, S. A., HALEY, B. K., CHENG, H. W., EINSTEIN, M. E. and**  
338 **RUBIN, D. A.** (2013) The effect of perch availability during pullet rearing and egg laying on  
339 musculoskeletal health of caged White Leghorn hens. *Poultry Science* **92**: 1972–1980.

340 **HUTCHINSON, J.R., ANDERSON, F. C., BLEMKER, S.S. and DELP, S. L.** (2005)  
341 Analysis of hindlimb muscle moment arms in *Tyrannosaurus rex* using a three-dimensional  
342 musculoskeletal computer model: implications for stance, gait, and speed. *Paleobiology* **31**: 676–  
343 701.

344 **HYRE, H. M.** (1955) The effect of heredity and environment on keel deformities in White  
345 Leghorns. *West Virginia Agricultural Experiment Station Bulletin.***381**.

346 **JACKSON, B. E. and DIAL, K. P.** (2011) Scaling of mechanical power output during burst  
347 escape flight in the Corvidae. *Journal of Experimental Biology* **214**: 452–461.

348 **KAPPELI, S., GEBHARDT-HENRICH, S. G., FROHLICH, E., PFULG, A. and**  
349 **STOFFEL, M. H.** (2011) Prevalence of keel bone deformities in Swiss laying hens. *British*  
350 *Poultry Science* **52**: 531–536.

351 **KONDO, K. L.** (2008) Osteoporotic vertebral compression fractures and vertebral augmentation.  
352 *Seminars in interventional radiology*. Thieme Medical Publishers pp. 413.

353 **KOZAK, M., TOBALSKE, B., MARTINS, C., WERBEL, H., and HARLANDER-**  
354 **MATAUSCHEK, A.** (2015) Chick- locomotion in a multilayer environment, accepted in *Poultry*  
355 *Science Association 104rd Annual Meeting*, Kentucky, USA.



356 **LEBLANC, C., TOBALSKE, B., WUERBEL, H. and HARLANDER-MATAUSCHEK, A.**  
 357 (2015) Locomotion skills of chicks over an inclined walkway. accepted in *Poultry Science*  
 358 *Association 104rd Annual Meeting*, Kentucky, USA.

359 **LIU, D., VEIT, H. P., WILSON, J. H. and DENBOW, D. M.** (2003) Long-term  
 360 supplementation of various dietary lipids alters bone mineral content, mechanical properties and  
 361 histological characteristics of Japanese quail. *Poultry Science* **82**: 831–839.

362 **LOPEZ, J.** (2007) Animal Welfare: Global Issues, Trends and Challenges. Scientific and  
 363 Technical Review, Vol. 24 (2). *Canadian. Veterinary. Journal.* **48**: 1163-1164.

364 **MOBERG, G. P.** (1985) Biological response to stress: Key to assessment of well-being, in:  
 365 MOBERG, G. P. (Ed) *American Physiological Society*, pp. 28-49 (Bethesda, MD).

366 **MOINARD, C., STATHAM, P. and GREEN, P. R.** (2004a) Control of landing flight by laying  
 367 hens: implications for the design of extensive housing systems. *British Poultry Science* **45**: 578–  
 368 584.

369 **MOINARD, C., STATHAM, P., HASKELL, M. J., MCCORQUODALE, C., JONES, R.B.**  
 370 **and GREEN, P. R.** (2004b) Accuracy of laying hens in jumping upwards and downwards  
 371 between perches in different light environments. *Applied Animal Behaviour Science* **85**: 77–92.

372 **NASR, M. A. F., MURELL, J. and NICOL, C. J.** (2013) The effect of keel fractures on egg  
 373 production, feed, and water consumption in individual laying hens. *British Poultry Science* **54**:  
 374 165–170.

375 **NASR, M. A. F., MURELL, J., WILKINGS, L. J. and NICOL, C. J.** (2012a) The effect of  
 376 two classes of opioid drug on the landing ability of laying hens with and without keel fractures.in  
 377 *UFAW Animal Welfare Conference: Recent Advances in Animal Welfare Science III*, York, UK.

378 **NASR, M. A. F., MURELL, J., WILKINGS, L. J. and NICOL, C. J.** (2012b) The effect of  
379 keel fractures on egg production parameters, mobility and behaviour in individual laying hens.  
380 *Animal Welfare* **21**: 127–135.

381 **NASR, M. A. F., MURELL, J., WILKINGS, L. J. and NICOL, C. J.** (2012c) Do Laying Hens  
382 with Keel Bone Fractures Experience Pain? *PLoS One* **7**:e42420.

383 **NASR, M. A. F., MURELL, J., WILKINGS, L. J. and NICOL, C. J.** (2015) The effects of  
384 two non-steroidal anti-inflammatory drugs on the mobility of laying hens with keel bone  
385 fractures. *Veterinary Anaesthesia and Analgesia***42**: 197-204.

386 **PETRIK, M. T., GUERIN, M. T. and WIDOWSKI, T. M.** (2014) On-farm comparison of keel  
387 fracture incidence in conventional cage and floor-housed laying hens. *Poultry Science*  
388 *Association 103rd Annual Meeting*, Corpus Christi, pp. 71

389 **PETRIK, M. T., GUERIN, M. T. and WIDOWSKI, T. M.** (2015) On-farm comparison of keel  
390 fracture prevalence and other welfare indicators in conventional cage and floor-housed laying  
391 hens in Ontario, Canada. Accepted to *Poultry Science*.

392 **PICKEL, T., SCSHOLZ, B. and SCHRADER, L.** (2010) Perch material and diameter affects  
393 particular perching behaviours in laying hens. *Applied Animal Behaviour Science* **127**: 37–42.

394 **PICKEL, T., SCSHOLZ, B. and SCHRADER, L.** (2011) Pressure load on keel bone and foot  
395 pads in perching laying hens in relation to perch design. *Poultry Science* **90**: 715–24.

396 **PRUNIER, A., MOUNIER, L., LE NEINDRE, P., LETERRIER, C., MORMÈDE,**  
397 **PAULMIER, V., PRUNET, P., TERLOUW, C. and GUATTEO, R.** (2013) Identifying and  
398 monitoring pain in farm animals: a review. *Animal* **7**: 998–1010.

399 **RATH, N. C., HUFF, G. R., HUFF, W. E. and BALOG, J. M.** (2000) Factors regulating bone  
400 maturity and strength in poultry. *Poultry Science* **79**: 1024–1032.

401 **RICHARDS, G. J., BROWN, S. N., BOOTH, F., TOSCANO, M. J. and WILKINS, L. J.**  
 402 (2012) Panic in free-range laying hens. *Veterinary Record* **170**: 519  
 403 **RODENBURG, T. B., TUYTTENS, F. A. M., DE REU, K., HERMAN, L., ZOONS, J. and SONCK, B.** (2008)  
 404 Welfare assessment of laying hens in furnished cages and non-cage systems : an on-farm  
 405 comparison. *Animal Welfare* **17**: 363–373.  
 406 **SANDILANDS, V., MOINARD, C. and SPARKS, N. H. C.** (2009) Providing laying hens with  
 407 perches: fulfilling behavioural needs but causing injury? *British Poultry Science* **50**: 395–406.  
 408 **SCHOLZ, B., KJAER, J. B. and SRADER, L.** (2014) Analysis of landing behaviour of three  
 409 layer lines on different perch designs. *British Poultry Science* **55**: 419-426.  
 410 **SCHOLZ, B., RÖNCHEN, S., HAMANN, H., HEWICKER-TRAUTWEIN, M. and DISTL,**  
 411 **O.** (2008) Keel bone condition in laying hens : a histological evaluation of macroscopically  
 412 assessed keel bones. *Berliner und Münchener Tierärztliche Wochenschrift* **121**: 89–94.  
 413 **SCOTT, G., LAMBE, N. R. and HITCHCOCK, D.** (1997) Ability of laying hens to negotiate  
 414 horizontal perches at different heights, separated by different angles. *British Poultry Science* **38**:  
 415 48–54.  
 416 **STRATMANN, A., FROHLICH, E. K. F., GEBHARDT-HENRICH, S., HARLANDER-**  
 417 **MATAUSCHEK, A., WÜRBEL, H. and TOSCANO, M. J.** (2015) Modification of aviary  
 418 design reduces incidence of falls, collisions and keel bone damage in laying hens. *Applied Animal*  
 419 *Behaviour Science* **165**: 112-123.  
 420 **STRATMANN, A., FROHLICH, E., WÜRBEL, H. and GEBHARDT-HENRICH, S. G.**  
 421 (2013) Crashes of laying hens in aviary systems. *Proceedings of the Joint Meeting of the 33rd*  
 422 *International Ethological Conference (IEC) & the Association for the Study of Animal Behaviour*  
 423 *(ASAB) Conference*, Newcastle-Gateshead, UK.

424 **STRATMANN, A., TOSCANO, M. J., FROHLICH, E. K. F., HARLANDER-**  
425 **MATAUSCHEK, A. and GEBHARDT-HENRICH, S.** Do soft perches reduce keel bone  
426 fractures in laying hens? Accepted to *PlosONE*.

427 **TARLTON, J. F., WILKINS, L. J., TOSCANO, M. J., AVERY, N. C. and KNOTT, L.**  
428 (2013) Reduced bone breakage and increased bone strength in free range laying hens fed omega-  
429 3 polyunsaturated fatty acid supplemented diets. *Bone* **52**: 578–586.

430 **THIRUVENKADAN, A. K., PANNEERSELVAM, S. and PRABAKARAN, R.** (2010) Layer  
431 breeding strategies: an overview. *Worlds Poultry Science Journal* **66**: 477–502.

432 **TIEMEIER, O. W.** (1941) Repaired bone injuries in birds. *Auk* **58**: 350–359.

433 **TILLMANN, J. E.** (2009) Fear of the dark: night-time roosting and anti-predation behaviour in  
434 the grey partridge (*Perdix perdix* L.). *Behaviour* **146**: 999–1023.

435 **TOBALSKE, B. W. and DIAL, K. P.** (2000) Effects of body size on take-off flight performance  
436 in the Phasianidae (Aves). *Journal of Experimental Biology* **203**: 3319–3332.

437 **TOBALSKE, B. W. and DIAL, K. P.** (2007) Aerodynamics of wing-assisted incline running in  
438 birds. *Journal of Experimental Biology* **210**: 1742–1751.

439 **TOBALSKE, B. W., HEDRICK, T. L., DIAL, K. P. and BIEWENER, A. A.** (2003)  
440 Comparative power curves in bird flight. *Nature* **421**: 363–366.

441 **TOMASZEWSKI, P. K., VERDONSCHOT, N., BULSTRA, S. K. and VERKERKE, G. J.**  
442 (2010) A comparative finite-element analysis of bone failure and load transfer of osseointegrated  
443 prostheses fixations. *Annals of Biomedical Engineering* **38**: 2418–2427.

444 **TOSCANO, M. J., BOOTH, F., WILKINS, L. J., AVERY, N. C., BROWN, S. B.,**  
445 **RICHARDS, G. and TARLTON, J. F.** The effects of long (C20/22) and short (C18) chain  
446 omega-3 fatty acids on keel bone fractures, bone biomechanics, behaviour and egg production in  
447 free range laying hens. Accepted to *Poultry Science*

448 **TOSCANO, M. J., BOOTH, F., WILKINS, L. J., BROWN, S. B., RICHARDS, G. and**  
449 **TARLTON, J. F.** (2014) Use of an impact tester to assess the likelihood of fractures occurring  
450 against key bird- and motion-related factors. *Proceedings of the 2014 Poultry Science*  
451 *Association Annual Meeting*, Corpus Christi.

452 **TOSCANO, M. J., WILKINS, L. J., MILBURN, G., THORPE, K. and TARLTON, J. F.**  
453 (2013) Development of an ex vivo protocol to model bone fracture in laying hens resulting from  
454 collisions. (PE Witten, Ed.). *PLoS One* **8**:e66215.

455 **WARREN, D. E.** (1937) Physiological and genetic studies of crooked keels in chickens. Kansas  
456 Agricultural Experiment Station Technical Bulletin:44.

457 **WATKINS, B. A., LI, Y., LIPPMAN, H. E. and FENG, S.** (2003) Modulatory effect of  
458 omega-3 polyunsaturated fatty acids on osteoblast function and bone metabolism. *Prostaglandins*  
459 *Leukotrienes Essential Fatty acids* **68**: 387–398.

460 **WHITEHEAD, C. C.** (2004a) Skeletal disorders in laying hens: the problem of osteoporosis and  
461 bone fractures. PERRY, G.C.. (Ed) *Welfare of the Laying Hen*, pp. 259-270 (Wallingford, CABI  
462 Publishing).

463 **WHITEHEAD, C. C.** (2004b) Overview of bone biology in the egg-laying hen. *Poultry Science*  
464 **83**: 193–199.

465 **WHITEHEAD, C. C. and FLEMING, R. H.** (2000) Osteoporosis in cage layers. *Poultry*  
466 *Science* **79**: 1033–1041.

467 **WILKINS, L. J., BROWN, S.N., ZIMMERMAN, P. H., LEEB, C. and NICOL, C. J.** (2004)  
468 Investigation of palpation as a method for determining the prevalence of keel and furculum  
469 damage in laying hens. *Veterinary Record* **155**: 547–549.

470    **WILKINS, L. J., MCKINSTRY, J. L., AVERY, N.C., KNOWLES, T. G., BROWN, S. N.,**  
471    **TARLTON, J and NICOL, C. J.** (2011) Influence of housing system and design on bone  
472    strength and keel bone fractures in laying hens. *Veteterinary Record* **169**: 414.